




PBH Update

10/17/2018

Step 1: Simulation

1. The mean number of PBHs in a time T' and V for R' is: $R'VT'$. Throw that many PBHs in V with your code. Save each one's r, ζ .
- Keep only the PBHs that are in HAWC's field of view (zenith angle $< 50^\circ$)




Step 2: Zebra

2. Calculate the number of counts from each PBH using:

(a) Calculate its average flux per unit time, area, and energy $F = \frac{1}{4\pi r^2 D} \frac{dN}{dE}$.

(b) Input F and ζ into Zebra to get a number of counts S

- Input flux and zenith angle into zebra to get the number of counts
- For now using only the lower part of the energy spectrum until spectrum from file option is added to zebra




Step 3: Make a histogram of the p-values

3. For each PBH, throw a value of $B(\zeta, D)$. This is the modeled background for your PBH
4. For each PBH, throw a “measured” value N by throwing with a Poissonian with mean $(S + B)$
5. For each PBH, determine the p-value p of N given the thrown B . Josh uses a Poisson valued as

$$\text{prob}(\geq N) = \sum_{i=N}^{\infty} \frac{B^i \exp(-B)}{i!} = 1 - \frac{\Gamma(N, B)}{\Gamma(N)} \quad (1.1)$$

where $\Gamma(N)$ is the gamma function and $\Gamma(N, B)$ is the upper incomplete gamma function.

Step 3: Make a histogram of the p-values



Next Steps

- Use more events
- Do the same for 1s, 10s, and 100s durations